

Table 5. Coexistence of HT with PTC or TC. Brief literature review.

Authors (reference)	Patients studied	HT patients with PTC n (%)
Dailey et al, 1955 (6)	278 with TC	35 (12.6)
Crile and Hazard, 1962 (7)	200 with HT	1 (0.5)
Ott et al, 1985 (8)	146 with HT and solitary cold nodules	47 (32)
Ott et al, 1987 (9)	161 with TC	61 (38)
Eisenberg et al, 1989 (10)	120 with TC	13 (10.8)
Sclafani et al, 1993 (11)	48 with HT	8 (17)
Schaffler et al, 1998 (12)	153 with TC	10 (6.5)
Loh et al, 1999 (13)	631 with DTC	128 (20.3)
Singh et al, 1999 (14)	388 with PTC	57 (15)
Kebebew et al, 2001 (15)	136 with PTC	41 (30)
Cipolla et al, 2005 (16)	71 with PTC	19 (26.7)
Labidi et al, 2006 (17)	78 with HT	11 (14.1)
Costanzo et al, 2006 (18)	282 thyroidectomized (46 DTC, 28 HT)	11 (3.9)
Larson et al, 2007 (19)	98 resected thyroid specimens with HT	43 (43.8)
Repplinger et al, 2008 (20)	1198 thyroidectomized (217 HT, 293 PTC)	63 (5.3)
Matesa-Anic et al, 2009 (21)	10508 undergoing FNAC	42 (0.4)
Mazokopakis et al, 2010	140 thyroidectomized (32 PTC, 42 HT)	12 (8.5)

HT: Hashimoto's Thyroiditis, TC: Thyroid Carcinoma, PTC: Papillary Thyroid Carcinoma, DTC: Differentiated Thyroid Carcinoma, FNAC: Fine Needle Aspiration Cytology