

**Table 1.** Comparisons between the women with functional hypoparathyroidism and those with secondary hyperparathyroidism

<b>Variable</b>	<b>Functional hypoparathyroidism (N=257)</b>	<b>Secondary hyperparathyroidism (N=148)</b>	<b>P</b>
Age (years), mean (SD)	78.7 (8.1)	82.4 (6.7)	<0.001
Body Mass Index (kg/m <sup>2</sup> ), mean (SD)	22.9 (4.1)	23.6 (4.2)	0.131
PTH (pg/ml)	47 (34-61)	107 (85-139)	<0.001
Phosphate (mg/dl)	3.7 (3.3-4.1)	3.4 (3.1-3.8)	<0.001
Albumin-adjusted calcium (mg/dl)	9.0 (8.2-9.5)	8.9 (7.8-9.5)	0.114
25-hydroxyvitamin D (ng/ml)	6.8 (3.8-9.0)	7.0 (4.0-8.9)	0.778
Estimated GFR (ml/min)	74.5 (63.8-87.8)	71.1 (56.0-84.3)	<0.001
Magnesium (mg/dl)	2.0 (1.9-2.1)	2.0 (2.0-2.1)	0.415
BMD (T-score), mean (SD)	-2.65 (0.83)	-2.88 (0.93)	0.010
BMD (g/cm <sup>2</sup> ), mean (SD)	0.55 (0.09)	0.52 (0.10)	0.010

Data are shown as median and interquartile range where not otherwise stated.