

**Table 1.** TSH and thyroid hormones concentration from TFT patients classified according to thyroid status categories

Category of TFT	TSH (mU/l)	TT4 ( $\mu\text{g/dl}$ )	T3 (ng/dl)
	Median (Q1-Q3)	mean $\pm$ SD	mean $\pm$ SD
Total (n=100)	1.7 (0.9-3.2)	7.7 $\pm$ 2.2	0.76 $\pm$ 0.34
OH (n=1)	0.2 (NA)	11 $\pm$ NA	0.98 $\pm$ NA
SH (n=7)	0.2 (0.04-0.3)	7.7 $\pm$ 2.2	1.07 $\pm$ 0.15
Eu (n=25)	1.5 (1.0-2.5)	8.6 $\pm$ 1.9	1.08 $\pm$ 0.26
oh (n=5)	8.8 (6.8-20.8)	4.16 $\pm$ 0.68	0.49 $\pm$ 0.21
sh (n=1)	8.6 (NA)	6.2 $\pm$ NA	1.1 $\pm$ NA
NTI (n=61)	1.7 (0.9-3.5)	7.54 $\pm$ 2.24	0.61 $\pm$ 0.28

OH and SH: overt and subclinical hyperthyroidism; oh and sh: overt and subclinical hypothyroidism; Eu: euthyroid; NTI: non-thyroidal illness; n: number of patients; NA: not available; SD: standard deviation.