

Research paper**Association of genetic variants of vit D binding protein (DBP/GC) and of the enzyme catalyzing its 25-hydroxylation (DCYP2R1) and serum vit D in postmenopausal women**

Wen Xu,¹ Jing Sun,¹ Wenbo Wang,² Xiran Wang,³ Yan Jiang,¹ Wei Huang,⁴ Xin Zheng,⁵ Qiuping Wang,⁶ Zhiwei Ning,⁷ Yu Pei,⁸ Min Nie,¹ Mei Li,¹ Ou Wang,¹ Xiaoping Xing,¹ Wei Yu,⁹ Qiang Lin,⁹ Ling Xu,¹⁰ Weibo Xia¹

¹Department of Endocrinology, Key Laboratory of Endocrinology, Ministry of Health, Peking Union Medical College Hospital, Chinese Academy of Medical Science; ²Department of Endocrinology, Peking University Shougang Hospital; ³Department of Cadre Unit, General Hospital of the second Artillery Force; ⁴Department of Endocrinology, BeiJing Haidian Hospital; ⁵Department of Endocrinology, China Rehabilitation Research Center, ⁶Department of Endocrinology, Beijing Liangxiang Hospital; ⁷Department of Endocrinology, Beijing Chaoyang Hospital; ⁸Department of Geriatric Endocrinology, Chinese PLA General Hospital; ⁹Department of Radiology, Peking Union Medical College Hospital, Chinese Academy of Medical Science; ¹⁰Department of Gynaecology and Obstetrics, Peking Union Medical College Hospital, Chinese Academy of Medical Science; Beijing, China

ABSTRACT

OBJECTIVE: To determine if GC (group-specific component globulin) and CYP2R1 genetic variants have an association with serum 25-OHD₃ levels, BMD or bone turnover markers in a population of Chinese postmenopausal women. **DESIGN:** We randomly selected 1494 postmenopausal women of the Han ethnic group from seven communities in Beijing. BMD was determined by dual energy X-ray absorptiometry; serum bone turnover markers and 25-OHD₃ were measured by the automated Roche electrochemiluminescence system; genotypes of GC and CYP2R1 were detected by the TaqMan allelic discrimination assay. Multiple statistic methods were used to test the associations of SNP genotypes and vitamin D levels. **RESULTS:** In our sample, 89.6% women had vitamin D deficiency and another 9.8% had vitamin D insufficiency. The variants of rs2298849 ($\beta=0.105$, $P<0.001$) in GC were significantly associated with serum 25-OHD₃ levels. Allele G of rs2298849 might be protective for serum 25-OHD₃ level. Among the haplotypes of rs222020-rs2298849, CG ($\beta=0.104$, $P=0.001$) corresponded to increasing serum 25-OHD₃ concentrations. CYP2R1 polymorphisms showed some significant association with serum β -CTX and PINP levels. **CONCLUSIONS:** We found that GC variants had a significant association with serum 25-OHD₃ levels among postmenopausal women of the Han ethnic group in Beijing, while CYP2R1 variants were not found to be significant.

Key words: Association analysis, BMD, Bone turnover markers, CYP2R1, GC, SNP, Vitamin D

Address for correspondence:

Prof. Weibo Xia, Department of Endocrinology, Key Laboratory of Endocrinology, Ministry of Health, Peking Union Medical College Hospital, Chinese Academy of Medical Science, Beijing 100730 China, Tel.: 86-10-6915-5358, Fax: 86-10-6915-5358, E-mail: xiaweibo@medmail.com.cn

Received 23-05-2013, Accepted 08-09-2013

INTRODUCTION

Vitamin D deficiency is a very common health problem. It is reported that worldwide more than 1 billion children and adults are at risk of vitamin D deficiency.¹ The situation is likely to be worse in China. According to a large sample (more than 3,000 individuals) cross-sectional study in older adults living in Beijing and Shanghai, the percentage of vitamin D deficiency and insufficiency were 69.2 and 24.4%, respectively.² Vitamin D deficiency may decrease peak bone mass in young people and increase the risk of osteoporosis, falls and fractures in the elderly and old.³ It also associated with increasing risk of several autoimmune, cardiovascular, infectious and metabolic diseases.¹ Although the main cause of vitamin D deficiency is considered to be lack of sunlight exposure and dietary vitamin D intake, family and twin studies have shown that genetic factors might also affect vitamin levels.^{4,6}

The genome-wide association study (GWAS) and other association studies have been carried out to determine the functional genes. Group-specific component globulin (GC), also called vitamin D binding protein (DBP), is a multifunctional protein which can bind to vitamin D, maintain the vitamin D level in the human body and transport vitamin D to target tissues for processing and utilization.⁷ CYP2R1 is a member of cytochrome P450 families. It can catalyze the 25-hydroxylation of vitamin D with no sex and species differences and shows catalytic activity toward both vitamin D₂ and D₃.⁸ Recently, some studies have observed an association of GC and CYP2R1 variants with serum vitamin D concentration in European populations.⁹⁻¹¹ However, it remains to be confirmed whether they have similar effects in Chinese populations. Based on the Peking Vertebral Fracture (PK-VF) study, which is a large-scale epidemiologic study with randomly selected postmenopausal women residing in communities in Beijing, we examined whether the association reported for GC (rs222020 and rs2298849) and CYP2R1 (rs12794714, rs10741657, rs1562902 and rs10766197) variants are also present in this sample. We also explored whether these variants have some relationship with bone mineral density (BMD) and bone turnover markers such as β -C-terminal telopeptide of type 1 collagen (β -CTX) and procollagen type I N propeptide (P1NP).

SUBJECTS AND METHODOLOGY

Participants

Our participants were from the PK-VF cohort together with randomly selected postmenopausal women living in communities in seven districts of Beijing. Before sampling, we put up posts on the bulletin board of each community center to explain the nature of the proposed study. We then made telephone calls to invite the randomly selected participants to join the study. Age distribution of our selected participants was consistent with the composition of those communities. At the end the PK-VF study, 2070 postmenopausal women were recruited. Every participant completed a questionnaire concerning items such as age, years since menopause, fracture history and medication history. Height and weight of the participants were measured and recorded by assistant doctors or nurses. Menopause was defined as the absence of menstruation for at least one year. Exclusion criteria were: 1) chronic liver disease; 2) chronic renal disease; 3) significant gastrointestinal diseases such as inflammatory bowel disease (IBD) and chronic diarrhea; 4) metabolic or inherited bone disease; 5) corticosteroid, anticonvulsant or anti-tuberculosis therapy for more than 6 months during the previous year. Finally, 1546 participants were included in our study. The study was approved by the Ethics Committee of the Peking Union Medical College Hospital (PUMCH). All participants signed informed consent forms before entering the study.

Biochemistry

We collected a fasting blood sample from all the participants. Serum concentrations of 25-hydroxyvitamin D₃ (25-OHD₃), β -CTX and P1NP were measured by a fully automated Roche electrochemiluminescence system (E170, Roche Diagnostics, Switzerland) at PUMCH. The detection limit of β -CTX, P1NP and 25-OHD₃ was 0.01 ng/ml, 5 ng/ml and 4 ng/ml, respectively. The intraassay and interassay CV were 2% and 3.1% for β -CTX, 2.3% and 1.7% for P1NP and 5.7% and 6.1% for 25-OHD₃.

BMD measurement

Details of BMD measurement have been described previously.¹² Briefly, we measured the BMD of the lumbar spine (L2-4), femoral neck (FN) and total hip by dual energy X-ray absorptiometry (DXA).

Genotyping

We searched for single nucleotide polymorphisms (SNP) information on the GC and CYP2R1 genes on the HapMap website (<http://hapmap.ncbi.nlm.nih.gov/>). SNPs were selected according to the following criteria: 1) high heterozygosity, which means minor allele frequency (MAF) higher than 20% in the Chinese population (CHB in HapMap); 2) classified as tag SNPs; 3) reported to be associated with serum 25-OHD₃ level in previous studies.⁹⁻¹¹ Finally, we selected two SNPs of the GC gene (rs222020, rs2298849) and four SNPs of the CYP2R1 gene (rs12794714, rs10741657, rs1562902, rs10766197). Each participant was genotyped for all six SNPs by TaqMan allelic discrimination assay (Applied Biosystems, USA). The whole reacting volume was 6µl, including 3µl (approximately 15 ng) sample DNA, 2.5µl TaqMan Universal PCR Master Mix, 0.0875µl TaqMan probe assay and 0.4125µl ddH₂O. Reactions were performed on a Real-Time PCR system of ABI Prism 7900 (Applied Biosystems, USA) in a 384-well reaction plate under standard conditions. The genotyping success rate for six SNPs was higher than 98%, and the concordance rate was about 94.4% based on 3% duplicated samples. Genotype frequencies of the six SNPs were similar to those in HapMap-CHB (<http://hapmap.ncbi.nlm.nih.gov/>) and they were all in Hardy-Weinberg equilibrium.

Statistics

The Hardy-Weinberg equilibrium (HWE) was tested using the goodness-of-fit chi-square test of the Haploview 4.2 program, which was also used to calculate normalized LD coefficient (*D'*), and the Pearson correlation coefficient (*r*²) to measure link-

age disequilibrium (LD) between pairwise SNPs. Haplotypes of each participant were determined by Phase 2.02. Serum 25-OHD₃ levels were square-root transformed to approximate normality. The independent samples t-test was performed to test the difference in serum 25-OHD₃ levels between different seasons. To establish the potential effect of numeric covariates on serum 25-OHD₃ levels, a Pearson correlation was conducted as an exploratory analysis (SPSS, version 17.0). Age, BMI and blood collecting season (winter-spring defined as 1, or summer defined as 2) were considered as potential covariates in a linear regression model. Association of the square-root-transformed serum 25-OHD₃ levels with age, season and all the SNPs and haplotypes were tested by general linear regression using backward elimination in SPSS, respectively. We also conducted the Pearson correlation or general linear regression to test correlation or association of the 6 SNPs with β-CTX, P1NP and BMD of L2-4, femoral neck and total hip. All analyses were carried out under an additive model. *P*<0.05 was considered statistically significant. The statistical power was calculated by Quanto software (<http://hydra.usc.edu/gxe/>). Our study achieved a power of more than 80%.

RESULTS

Basic characteristics of participants

After excluding participants who failed the biochemistry test or genotyping, 1494 participants remained. The major clinical profiles of these participants are listed in Table 1. Among them, 1338 women had vitamin D deficiency (25-OHD₃ <20ng/ml) and another 146 women had vitamin D insuffi-

Table 1. Basic characteristics of participants

Characteristics	Mean ± SD	Characteristics	Mean ± SD
Age (years) (n=1494)	64.0 ± 9.2	Vitamin D deficiency (n, %)	1338, 89.6%
Height (cm) (n=1494)	155.1 ± 5.6	Vitamin D insufficiency (n, %)	146, 9.8%
Weight (kg) (n=1494)	62.8 ± 9.8	β-CTX (ng/ml) (n=1494)	0.440 ± 0.212
Body mass index (kg/m ²) (n=1494)	26.1 ± 3.7	P1NP (ng/ml) (n=1494)	56.95 ± 28.70
Serum 25-OHD ₃ (ng/ml) (n=1494)	13.2 ± 5.4	BMD of L2-4 (g/cm ²) (n=1452)	0.966 ± 0.198
Winter-Spring (Dec-May) (n=805)	12.8 ± 5.2	BMD of FN (g/cm ²) (n=1451)	0.767 ± 0.139
Summer (Jun-Aug) (n=689)	13.7 ± 5.6	BMD of TH (g/cm ²) (n=985)	0.865 ± 0.144

Vitamin D deficiency: 25-OHD₃ <20ng/ml; vitamin D insufficiency: 20ng/ml ≤ 25-OHD₃ <30ng/ml.

ciency (20ng/ml \leq 25-OHD₃ <30ng/ml), accounting for 89.6% and 9.8% of the total sample, respectively.

Among the 1494 participants, 91 blood samples were collected in winter, while 714 and 689 blood samples were collected in spring and summer, respectively. Since the sample size of the winter group was small and there was no significantly statistical difference between the winter and spring groups after adjusting age and BMI, we divided the 1494 participants into two season groups: summer and winter-spring. The concentrations of serum 25-OHD₃ were significantly different between the summer group and the winter-spring group ($P=0.001$).

Serum 25-OHD₃ levels in different seasons

Because the sample size of the winter group was small, and there was no significantly statistical difference of serum 25-OHD₃ levels between the winter and spring groups ($P=0.671$), we divided the 1494 participants into two season groups: summer and winter-spring. The serum 25-OHD₃ levels were significantly different between the summer and the winter-spring groups ($P=0.002$), as shown in Table 2.

Association of GC and CYP2R1 genetic variants with serum 25-OHD₃ levels

To determine the potential effect of numeric covariates on serum 25-OHD₃ levels, a Pearson correlation was conducted as an exploratory analysis (SPSS, version 17.0). Except for all the SNPs under the additive model, age, BMI and blood collecting

Table 2. Result of t-test of serum 25-OHD₃ levels in different seasons

Blood collecting season	N	25-OHD ₃ (ng/ml)	Sqrt of 25-OHD ₃ (ng/ml)	P
Winter	91	13.1 ± 5.7	3.54 ± 0.80	0.671
Spring	714	12.8 ± 5.1	3.50 ± 0.72	
Winter - Spring (Dec-May)	805	12.8 ± 5.2	3.51 ± 0.73	
Summer (Jun-Aug)	689	13.7 ± 5.6	3.62 ± 0.77	0.002

Sqrt of 25-OHD₃: square-root-transformed serum 25-OHD₃ level.

season (winter-spring defined as 1, or summer defined as 2) were considered as potential covariates in a linear regression model. In our study, SNPs rs222020 ($r=0.080$, $P=0.001$) and rs2298849 ($r=0.095$, $P<0.001$) of the GC gene, age ($r=-0.074$, $P=0.002$) and blood collecting season ($r=0.079$, $P=0.001$) were found to have significant correlation with serum 25-OHD₃ levels. Following this, the association of the square-root-transformed serum 25-OHD₃ levels with age, season and all the SNPs were tested by general linear regression using backward elimination in SPSS. The main results of the SNPs genotyping of the GC and the CYP2R1 genes and their association analyses for serum 25-OHD₃ levels are shown in Table 3. SNP rs2298849 ($\beta=0.105$, $P<0.001$) of the GC gene was found to have a significant association with serum 25-OHD₃ levels. We did not find any significant association between CYP2R1 genetic variants and serum 25-OHD₃ levels.

Table 3. Results of single SNPs association analyses for serum 25-OHD₃ levels

Covariates					r	PI	β (SE)	P2
Genes	SNP	Allele	MAF	HWE				
GC	rs222020	T/C	0.338	0.86	0.080	0.001		
GC	rs2298849	A/G	0.324	0.55	0.095	<0.001	0.105(0.029)	<0.001
CYP2R1	rs12794714	G/A	0.373	0.87	-0.010	0.353		
CYP2R1	rs10741657	G/A	0.396	0.96	-0.007	0.387		
CYP2R1	rs1562902	T/C	0.435	0.83	-0.016	0.272		
CYP2R1	rs10766197	G/A	0.371	0.51	-0.014	0.296		
Age					-0.074	0.002	-0.006(0.002)	0.003
Season					0.079	0.001	0.127(0.039)	0.001

Serum 25-OHD₃ levels were square-root transformed to approximate normality. Bold numbers represent significant P values. Allele: major allele/minor allele; MAF: minor allele frequency; HWE: P values for Hardy-Weinberg Equilibrium test; r: Pearson correlation coefficient; PI: P value of Pearson correlation; β : regression coefficient; SE: standard error; P2: P value of general linear regression.

The LD plots of the GC and CYP2R1 genes are depicted in Figure 1. The results of the haplotype association analyses are displayed in Table 4. The two SNPs of the GC gene were in relatively strong




Figure 1. LD plots with D' values of the GC (a) and CYP2R1 (b) genes in our 1494 participants. The Figure was generated by Haploview. Block definition followed the rules of solid spine of LD. D' values multiplied by 100 are shown as a number in the diamonds.

Table 4. Results of haplotypes association analyses for serum 25-OHD₃ levels

Covariates	r	PI	β (SE)	P2
GC: rs222020-rs2298849				
Haplotype	Frequency			
TA	0.638	-0.082	0.001	
CG	0.299	0.094	< 0.001	0.104 (0.030) 0.001
CA	0.038	-0.024	0.173	
TG	0.024	0.012	0.322	
CYP2R1: rs12794714-rs10741657-rs1562902-rs10766197				
Haplotype	Frequency			
GACG	0.375	-0.004	0.444	
AGTA	0.347	-0.019	0.233	
GGTG	0.165	0.030	0.124	
GGCG	0.057	-0.017	0.250	
Age		-0.074	0.002	-0.006 (0.002) 0.003
Season		0.079	0.001	0.125 (0.039) 0.001

Serum 25-OHD₃ levels were square-root transformed to approximate normality. Bold numbers represent significant P values. r: Pearson correlation coefficient; PI: P value of Pearson correlation; β : regression coefficient; SE: standard error; P2: P value of general linear regression.

LD ($D' > 0.8$) with each other, as were also the four SNPs of the CYP2R1 gene. P values were calculated using general linear regression (SPSS, version 17.0) under an additive model for those haplotypes with more than 1% frequency. Among the haplotypes of rs222020-rs2298849, CG ($\beta = 0.104$, $P = 0.001$) corresponded to increasing serum 25-OHD₃ concentrations. We did not find any significant association between haplotypes of the CYP2R1 gene and serum 25-OHD₃ levels.

Association of GC and CYP2R1 genetic variants with bone turnover markers and BMD

Basic information concerning P1NP, β -CTX and BMD of the lumbar spine (L2-4), femoral neck (FN) and total hip (TH) is shown in Table 1. Some participants failed to attend the BMD test, therefore the available data of BMD was limited, especially as regards BMD of TH.

To determine the potential effect of numeric covariates on bone turnover markers and BMD, a Pearson correlation was conducted (SPSS, version 17.0) as an exploratory analysis. Age and BMI were considered as potential covariates.

As shown in Table 5, in our research SNP rs12794714 ($r = -0.053$, $P = 0.021$) and BMI ($r = -0.052$, $P = 0.022$) were found to have significant correlation with β -CTX levels; SNPs rs12794714 ($r = -0.064$, $P = 0.007$) and rs10741657 ($r = 0.046$, $P = 0.036$) had significant correlation with P1NP levels. In the final linear regression model, SNPs rs12794714 ($\beta = -0.056$, $P = 0.002$) and rs10766197 ($\beta = 0.044$, $P = 0.012$) had statistically significant association with β -CTX levels; SNP rs12794714 ($\beta = -0.017$, $P = 0.014$) had statistically significant association with P1NP levels. Although we found some statistically significant results, as the magnitude of association was rather small, we did not do further haplotype association analysis.

As shown in Table 6, in our research, age ($r = -0.299$, $P < 0.001$) and BMI ($r = 0.183$, $P < 0.001$) had significant correlation with BMD of L2-4; age ($r = -0.521$, $P < 0.001$) and BMI ($r = 0.210$, $P < 0.001$) had significant correlation with BMD of FN; age ($r = -0.535$, $P < 0.001$) and BMI ($r = 0.272$, $P < 0.001$) had significant correlation with BMD of TH. None of the 6 SNPs were found to have significant correlation with BMD of

Table 5. Results of single SNPs association analyses for β -CTX and P1NP levels

Dependent Covariates		β -CTX				P1NP			
		r	PI	β (SE)	P2	r	PI	β (SE)	P2
Gene	SNP								
GC	rs222020	0.010	0.349			0.008	0.379		
GC	rs2298849	-0.004	0.436			-0.003	0.457		
CYP2R1	rs12794714	-0.053	0.021	-0.056 (0.018)	0.002	-0.064	0.007	-0.017 (0.007)	0.014
CYP2R1	rs10741657	0.036	0.084			0.046	0.036		
CYP2R1	rs1562902	0.025	0.163			0.036	0.085		
CYP2R1	rs10766197	-0.018	0.246	0.044 (0.018)	0.012	-0.043	0.050		
Age		0.029	0.128			0.030	0.121		
BMI		-0.052	0.022	-0.003 (0.001)	0.051	0.010	0.351		

P1NP levels were log-transformed to approximate normality. Bold numbers represent significant *P* values.

r: Pearson correlation coefficient; PI: *P* value of Pearson correlation; β : regression coefficient; SE: standard error; P2: *P* value of general linear regression.

Table 6. Results of single SNPs correlation analyses for BMD of L2-4, FN, and TH

Dependent Covariates		L2-4 BMD (N=1452)		FN BMD (N=1451)		TH BMD (N=985)	
		r	<i>P</i>	r	<i>P</i>	r	<i>P</i>
Gene	SNP						
GC	rs222020	-0.019	0.231	0.014	0.291	-0.006	0.430
GC	rs2298849	-0.019	0.237	0.009	0.361	-0.013	0.347
CYP2R1	rs12794714	0.005	0.431	0.017	0.261	0.031	0.163
CYP2R1	rs10741657	0.003	0.452	-0.014	0.301	0.001	0.491
CYP2R1	rs1562902	0.001	0.492	-0.025	0.173	-0.033	0.149
CYP2R1	rs10766197	-0.002	0.463	0.011	0.344	0.038	0.118
Age		-0.299	<0.001	-0.521	<0.001	-0.535	<0.001
BMI		0.183	<0.001	0.210	<0.001	0.272	<0.001

Bold numbers represent significant *P* values. r: Pearson correlation coefficient; *P*: *P* value of Pearson correlation.

L2-4, FN, or TH ($P > 0.05$). We thus did not proceed to further linear regression to test the association between SNPs or haplotypes and BMD.

DISCUSSION

In our study, 89.6% of the 1494 women had vitamin D deficiency, which suggests that vitamin D deficiency might be very common in postmenopausal women in Beijing. percentage is higher than the 69.2% reported in a previous study by Lu et al. in a Chinese population.² The observed difference might be due to age, latitude, sex and blood collecting season. The average age of our sample was about 64 years old,

which is higher than the 59 years old of their study. Participants of our study were residents of Beijing, while theirs lived in Beijing and Shanghai and, as is known, Beijing is at higher latitude than Shanghai. Our participants were all women, while they included both men and women. They collected blood samples mainly from April to June, which is characterized by relatively abundant sunlight exposure, while we collected blood samples over a large scale of time with less sunlight, from December to October.

We found that the variants of rs2298849 ($\beta = 0.105$, $P < 0.001$) in the GC gene were significantly associated with serum 25-OHD₃ levels. Allele G of rs2298849

might be protective for serum 25-OHD₃ levels. The variants of rs222020 ($r=0.080$, $P=0.001$) in the GC gene were correlated with serum 25-OHD₃ levels, although they were not significantly associated with serum 25-OHD₃ levels in the final linear regression model. This finding is consistent with the study by Bu et al.¹¹ Previous studies on the GC gene focused more on functional SNP rs4588 located in the exon. Its A/Lyn allele corresponds to the GC2 protein, which was found to be significantly associated with decreased serum 25-OHD₃ level^{7,14,15} Lu et al. observed that the variants of rs4588 were associated with plasma 25-OHD₃ levels in a Chinese population,¹⁶ while Ahn et al. and Wang et al. found that rs2282679 variants were associated with serum 25-OHD₃ levels in white Europeans.^{9,10} In the HapMap Data, rs2282679 and rs4588 are in complete LD ($r^2=1$) in CEU and CHB populations, while these two SNPs are both in very weak LD with rs222020 and rs2298849 (r^2 ranging from 0.08 to 0.15). As rs222020 and rs2298849 are both located in the intron, there might be some new functional site which is independent of rs4588 and can influence serum 25-OHD₃ levels. Fine genotyping in the area near rs222020 and rs2298849 might need to be done to discover new functional sites.

We detected no significant association between CYP2R1 polymorphisms and serum 25-OHD₃ levels in Chinese postmenopausal women. This finding is consistent with the previous study in a Chinese population by Lu et al.¹⁶ However, CYP2R1 genetic variants were found to be associated with vitamin D level in many studies in white Europeans.⁹⁻¹¹ Since we had >99% power to find the beta values reported by Bu et al. in white Europeans,¹¹ the discrepancy with previous studies may result from LD pattern variation of the causal SNP and the four SNPs detected in our study between Chinese Hans and white Europeans. Previous studies have hypothesized that genetic factors may play a much more important role in influencing vitamin D level in men than in women.⁴ As we had only female participants, our results may not represent the status of men. Another possibility is that other vitamin D 25-hydroxylation enzymes play a more important role in Chinese Hans than in white Europeans, thus CYP2R1 polymorphisms would hypothetically not influence serum 25-OHD₃ level in Chinese Hans.

In our study, we found CYP2R1 polymorphisms to be significantly associated with two bone turnover markers: β -CTX and P1NP. As we discovered no significant association between CYP2R1 polymorphisms and serum 25-OHD₃ levels in our study, the CYP2R1 gene might influence these two bone turnover markers via some as yet unknown mechanism. Since we did not locate any previous studies on the association between CYP2R1 genetic polymorphisms and the two bone turnover markers, this result may need to be further corroborated considering the small magnitude of the association.

CONCLUSIONS

In conclusion, we found that GC genetic variants had a significant association with serum 25-OHD₃ level among postmenopausal women of the Han ethnic group in Beijing, while CYP2R1 genetic variants were not observed as being significant. Association of CYP2R1 genetic variants with β -CTX and P1NP needs to be further substantiated.

GRANTS SUPPORTS

This research was supported by the National Natural Science Foundation of China (No.81070687 and 81170805), National Science and Technology Pillar Program (2006BAI02B03), National Science and Technology Major Projects for "Major New Drugs Innovation and Development" (Grant2008ZX09312-016), Beijing Natural Science Foundation (No. 7121012) and Scientific Research Foundation of Beijing Medical Development (No. 2007-3029).

REFERENCES

1. Holick MF, 2011 Vitamin D deficiency in 2010: health benefits of vitamin D and sunlight: a D-bate. *Nat Rev Endocrinol* 7: 73-75.
2. Lu L, Yu Z, Pan A, et al, 2009 Plasma 25-hydroxyvitamin D concentration and metabolic syndrome among middle-aged and elderly Chinese individuals. *Diabetes Care* 32: 1278-1283.
3. Holick MF, 2011 Vitamin D: a d-lightful solution for health. *J Investig Med* 59: 872-880.
4. Arguelles LM, Langman CB, Ariza AJ, et al, 2009 Heritability and environmental factors affecting vitamin D status in rural Chinese adolescent twins. *J Clin Endocrinol Metab* 94: 3273-3281.

5. Shea MK, Benjamin EJ, Dupuis J, et al, 2009 Genetic and non-genetic correlates of vitamins K and D. *Eur J Clin Nutr* 63: 458-464.
6. Snellman G, Melhus H, Gedeberg R, et al, 2009 Seasonal genetic influence on serum 25-hydroxyvitamin D levels: a twin study. *PLoS One* 4: e7747.
7. Fang Y, van Meurs JB, Arp P, et al, 2009 Vitamin D binding protein genotype and osteoporosis. *Calcif Tissue Int* 85: 85-93.
8. Strushkevich N, Usanov SA, Plotnikov AN, Jones G, Park HW, 2008 Structural analysis of CYP2R1 in complex with vitamin D3. *J Mol Biol* 380: 95-106.
9. Ahn J, Yu K, Stolzenberg-Solomon R, et al, 2010 Genome-wide association study of circulating vitamin D levels. *Hum Mol Genet* 19: 2739-2745.
10. Wang TJ, Zhang F, Richards JB, et al, 2010 Common genetic determinants of vitamin D insufficiency: a genome-wide association study. *Lancet* 376: 180-188.
11. Bu FX, Armas L, Lappe J, et al, 2010 Comprehensive association analysis of nine candidate genes with serum 25-hydroxy vitamin D levels among healthy Caucasian subjects. *Hum Genet* 128: 549-56.
12. Zhao J, Xia W, Nie M, et al, 2012 A haplotype of MATN3 is associated with vertebral fracture in Chinese postmenopausal women: Peking Vertebral Fracture (PK-VF) study. *Bone* 50: 917-924.
13. Purcell S, Neale B, Todd-Brown K, et al, 2007 PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81: 559-575.
14. Abbas S, Linseisen J, Slinger T, et al, 2008 The Gc2 allele of the vitamin D binding protein is associated with a decreased postmenopausal breast cancer risk, independent of the vitamin D status. *Cancer Epidemiol Biomarkers Prev* 17: 1339-1343.
15. Lauridsen AL, Vestergaard P, Hermann AP, et al, 2005 Plasma concentrations of 25-hydroxy-vitamin D and 1,25-dihydroxy-vitamin D are related to the phenotype of Gc (vitamin D-binding protein): a cross-sectional study on 595 early postmenopausal women. *Calcif Tissue Int* 77: 15-22.
16. Lu L, Sheng H, Li H, et al, 2012 Associations between common variants in GC and DHCR7/NADSYN1 and vitamin D concentration in Chinese Hans. *Hum Genet* 131: 505-512.